

Spreading Spondyloarthritis: are ILCs cytokine shuttles from base camp gut?

Barbara Neerinckx^{1,2}, Dirk Elewaut^{3,4}, Rik J. Lories^{1,2}

¹Laboratory of Tissue Homeostasis and Disease, Skeletal Biology and Engineering Research Center, KU Leuven; ² Division of Rheumatology, University Hospitals Leuven; ³ Laboratory of Molecular Immunology and Inflammation, VIB Inflammation Research Center, Ghent University, Belgium⁴ Division of Rheumatology, University of Gent, Belgium.

Corresponding author: Rik Lories, SBE Research Center, O&N1, Box 813, B3000 Leuven, Belgium.
Phone: +32-16-342541; Fax: +32-16-342543; e-mail: Rik.Lories@uz.kuleuven.be

Conflict of interest: The authors have no conflict of interest. Collaborative research of the authors on the topic is supported by grant G.0946.14 from the Flanders Research Foundation and by the DevRepair (P7/07) IAP-VII network (Belgian Federal Government).

A series of discoveries has transformed concepts of spondyloarthritis and pro-inflammatory cytokine interleukin-23 (IL-23) has taken center stage. The IL-23 frenzy kicked off with the identification of SNPs in the IL-23 receptor (IL23R) gene that are associated with ankylosing spondylitis (AS) ¹ and also with related disorders such as psoriasis, psoriatic arthritis and Crohn's disease. The most striking and direct evidence comes from an *in vivo* IL-23 overexpression study in mice that phenocopies the human disease and its Janus-faced characteristics: joint inflammation and structural damage presenting as new bone formation². Patient studies reported increased serum levels of IL-23 in AS ³⁻⁵ and presence of IL-23 positive cells was shown in facet joints of AS patients⁶. Direct clinical evidence comes from a prospective, open-label clinical trial with ustekinumab, an antibody binding to the shared p40 subunit of IL-23 and IL-12 ⁷ and successful clinical trials targeting IL-17, one of the downstream cytokines associated with IL-23 signaling⁸. Key to the hypothesis and evidence proposed in the mouse model, is the presence of an IL-23 receptor positive T-cell population in the entheses of mice. As enthesitis is one of the main characteristics of AS and was proposed as the primary lesion⁹, it is hard to ignore the potential key role of such cells. However, until now, these IL-23 receptor positive cells have not yet been demonstrated in human samples.

In this issue of ARD, *Ciccia* and colleagues report on the presence of a population of IL-23R positive innate lymphoid cells (ILCs) in the gut, peripheral blood, synovial fluid and bone marrow of AS patients¹⁰. Numbers of such cells are increased as compared to different controls and their surface characteristics show similarities with the mouse cells identified earlier. However, these ILCs are a rare cell type whose significance in human disease is as yet uncertain. In two previous studies *Ciccia et al* showed increased IL-23 expression in the gut of AS patients as compared to healthy controls^{11,12}. In AS synovium and peripheral blood cells however, no differences in expression of IL-23 were found^{13,14}.

They characterized these cells as Lyn⁻IL-23R⁺NKp44⁺Tbet⁺RORc⁻ cells and their induced production of cytokines IL-17 and IL-22 led them to conclude these are ILC3 cells. Unlike conventional ILC3 and IL-23R⁺ enthesal T cells, these cells seemingly do not express RORc but do express the transcription factor Tbet which can, according to the authors, possibly be explained by a specific stage of differentiation of these cells. This finding makes them somewhat atypical, even for ILC3s. The reasons of these differences between 'conventional' characteristics of ILCs and the spondyloarthritis' associated cells are unclear and whether these changes precede onset of inflammation or are rather a secondary to already established inflammation remains to be determined. Further clarification of

the exact nature of these cells clearly needs to be carried out. Hence, fine details in small subpopulations of cells but may still represent a static rather than dynamic view on cell populations that are actively part of host defense or disease.

The expansion of these ILC3 cells in AS patients with acute and chronic gut inflammation was significantly correlated with the disease activity as assessed by the BASDAI. In AS patients without gut inflammation the upregulation of ILC3s was not detected. The authors not only reported an expansion of NKp44+ILC3s in gut samples of AS patients, but also in the peripheral blood, synovial fluid and the bone marrow of AS patients. ILC3s are defined by their capacity to produce the interleukins IL-17 and IL-22¹⁵, portrayed as critical cytokines in the pathogenesis of AS. In the study, gut ILC3 cells were demonstrated to produce IL-17 and IL-22 and a small percentage of cells expressed both cytokines. In the peripheral blood, the majority of ILC3s produced IL-22 and only a small subset of ILC3s produced IL-17 or the combination of IL-22 and IL-17. Among synovial fluid and bone marrow mononuclear ILC3s produced exclusively IL-22. The reasons for these differential patterns according to localization are unclear and could reflect tissue imprinted cytokine patterns. These data suggest ILCs could be an important supplier of IL-17 and IL-22 in AS, which is also one of the implicit conclusions of this study by *Ciccia et al.*

Insights into the role of IL-17 in spondyloarthritis are dynamically evolving with a shift from a focus on adaptive immune cells towards more innate immune populations. Indeed, IL-17 producing CD4+ cells, also known as Th17 cells, were shown to be elevated in HLA-B27/human beta2m transgenic rats¹⁶ and in the lymph nodes of an AS mouse model¹⁷. Moreover, there are studies demonstrating an increased amount of IL-17 producing CD4+ cells in the blood of AS patients as compared to controls¹⁸⁻²¹ albeit that this observation is not consistent²²⁻²⁴. IL-17 producing cells were also found in the subchondral bone marrow cells of affected facet joints of AS patients⁶. Interestingly, the majority of these IL17 positive cells were rather innate immune cells (CD15+ neutrophils and MPO+ cells of the myeloid lineage) than CD4+ T cells⁶. Obviously, the most convincing proof of principle that IL-17 plays a key role in spondyloarthritis is found in the clinical trials evaluating the effect of secukinumab, a monoclonal anti-IL17A antibody that rapidly reduced clinical and biological signs of active ankylosing spondylitis⁸.

IL-22 has also earned its place in the spotlights as it has not only been associated with enthesitis/arthritis development in mice but also with upregulation of genes potentially involved in the new bone formation process (such as Wnts and bone morphogenetic proteins)². However, the role for IL-22 in AS and related disorders seems to be tissue dependent - as demonstrated in a study by *Benham et al* using the SKG mouse model²⁵. In this mouse model curdlan injection in SKG mice leads to IL-23 dependent axial and peripheral arthritis and ileitis. When IL-22 was neutralized by an IL-22 antibody, the mice developed a reduced severity of enthesitis but an exacerbation of ileitis²⁵. This finding highlights that the effects of interleukins in the pathogenesis of AS are not always straight forward, but can vary from tissue to tissue and influenced by tissue dependent environmental factors. It is important to note that the current attention towards IL-23, IL-17 and IL-22 somewhat neglects the critical role of tumor necrosis factor (TNF) in this type of diseases. Not only does a TNF overexpression model mimic many features of the disease²⁶ but TNF inhibition is currently the mainstay approach with high and sustained efficacy in SpA. Bringing together the different proinflammatory cytokines in one paradigm is therefore an interesting challenge.

Detailed histological analysis of the gut biopsies drives the authors to propose the hypothesis that IL-23 receptor positive ILC3s differentiate in the gut and then migrate to extra-intestinal sites where they produce IL-17 and IL-22. This hypothesis was formulated in part on the abundant expression of $\alpha 4\beta 7$ integrin on ILC3s, a $\beta 7$ integrin with marked gut tropism, even though earlier work demonstrated its presence on synovial T cells in SpA synovium²⁷. Although this hypothesis touches the limits of careful observation in human samples without additional mechanistic experiments, the

concept is certainly worth considering. Moreover, earlier this year *Mackley et al* provided the first *in vivo* evidence of ILC3 traffic starting from the gut in mice²⁸. The concept is also in line with old and new insights into the role of gut inflammation in spondyloarthritis. Five to 10% of AS patients suffer from clinically apparent inflammatory bowel disease (IBD) and even a much larger proportion of AS patients suffer from subclinical gut inflammation as already suggested in 1985²⁹. In a recent study by *Van Praet et al* 46.2% of AS patients showed microscopic gut inflammation of which 16.9% was acute and 29.2% was chronic inflammation³⁰. AS patients with chronic gut inflammation also have a higher degree of bone marrow edema, as seen on MRI further supporting a link between mucosal inflammation and progressive disease in AS³¹. The gut is a barrier tissue continuously faced with the challenge of maintaining the symbiotic gut microbiome while avoiding local and systemic disease development by pathogenic microorganisms. New technologies can assess the gut microbiome in a systematic way. Different efforts are underway to link the microbiome with gut inflammation and joint disease in SpA and the results are eagerly awaited.

ILC3s are required for human immunity against extracellular bacteria and also play a role in chronic inflammation due to its pro-inflammatory characteristics¹⁵. Keeping in mind that intestinal bacteria could play a role in the pathogenesis of AS, as described above, and that AS is a chronic inflammatory disease, the hypothesis that ILC3s are directly involved in the pathogenesis of AS sounds plausible. Of course, further research is necessary to, firstly, confirm the data and, secondly, further explore this hypothesis. If the hypothesis about the migration of IL-23 sensitized gut-resident ILC3s is correct, one of the key residual questions for example is how HLA-B27, the strongest risk factor for AS, fits in this story. Since HLA-B27 is an MCH class I molecule it will present endogenous antigens/peptides originating from the cytoplasm on the cell surface. These antigens/peptides do not only include self-peptides but can also derive from viruses and bacteria. This peptide presentation could then possibly be the link to the gut microbiome. As the HLA-B27 gene is not linked to IBD, another possibility is that HLA-B27 does not play a role in disease induction in the gut, but only plays a role in inflammation at the joint level. Another question is how to explain joint inflammation in AS patients without currently detectable gut inflammation. Notwithstanding the many additional questions, the paradigm proposed by Ciccia et al focusing on ILC3 cells as cytokine shuttle travelling from gut to blood, synovium and bone marrow in AS is an interesting new approach to the unraveling of spondyloarthritis' pathogenesis.

- 1 Burton, P. R. *et al*. Association scan of 14,500 nonsynonymous SNPs in four diseases identifies autoimmunity variants. *Nature genetics* **39**, 1329-1337, doi:10.1038/ng.2007.17 (2007).
- 2 Sherlock, J. P. *et al*. IL-23 induces spondyloarthropathy by acting on ROR-gammat+ CD3+CD4- CD8- enthesal resident T cells. *Nature medicine* **18**, 1069-1076, doi:10.1038/nm.2817 (2012).
- 3 Wang, X., Lin, Z., Wei, Q., Jiang, Y. & Gu, J. Expression of IL-23 and IL-17 and effect of IL-23 on IL-17 production in ankylosing spondylitis. *Rheumatology international* **29**, 1343-1347, doi:10.1007/s00296-009-0883-x (2009).
- 4 Mei, Y. *et al*. Increased serum IL-17 and IL-23 in the patient with ankylosing spondylitis. *Clinical rheumatology* **30**, 269-273, doi:10.1007/s10067-010-1647-4 (2011).
- 5 Andersen, T. *et al*. Increased plasma levels of IL-21 and IL-23 in spondyloarthritis are not associated with clinical and MRI findings. *Rheumatology international* **32**, 387-393, doi:10.1007/s00296-010-1655-3 (2012).
- 6 Appel, H. *et al*. In situ analysis of interleukin-23- and interleukin-12-positive cells in the spine of patients with ankylosing spondylitis. *Arthritis and rheumatism* **65**, 1522-1529, doi:10.1002/art.37937 (2013).

- 7 Poddubnyy, D., Hermann, K. G., Callhoff, J., Listing, J. & Sieper, J. Ustekinumab for the treatment of patients with active ankylosing spondylitis: results of a 28-week, prospective, open-label, proof-of-concept study (TOPAS). *Annals of the rheumatic diseases* **73**, 817-823, doi:10.1136/annrheumdis-2013-204248 (2014).
- 8 Baeten, D. *et al.* Anti-interleukin-17A monoclonal antibody secukinumab in treatment of ankylosing spondylitis: a randomised, double-blind, placebo-controlled trial. *Lancet* **382**, 1705-1713, doi:10.1016/s0140-6736(13)61134-4 (2013).
- 9 Benjamin, M. & McGonagle, D. The anatomical basis for disease localisation in seronegative spondyloarthropathy at entheses and related sites. *Journal of anatomy* **199**, 503-526 (2001).
- 10 Ciccia, F. *et al.* Type 3 innate lymphoid cells producing IL-17 and IL-22 are expanded in the gut, in the peripheral blood, synovial fluid and bone marrow of patients with ankylosing spondylitis. *Annals of the rheumatic diseases*, doi:10.1136/annrheumdis-2014-206323 (2015).
- 11 Ciccia, F. *et al.* Overexpression of interleukin-23, but not interleukin-17, as an immunologic signature of subclinical intestinal inflammation in ankylosing spondylitis. *Arthritis and rheumatism* **60**, 955-965, doi:10.1002/art.24389 (2009).
- 12 Ciccia, F. *et al.* Evidence that autophagy, but not the unfolded protein response, regulates the expression of IL-23 in the gut of patients with ankylosing spondylitis and subclinical gut inflammation. *Annals of the rheumatic diseases* **73**, 1566-1574, doi:10.1136/annrheumdis-2012-202925 (2014).
- 13 Neerinckx, B., Carter, S. & Lories, R. IL-23 expression and activation of autophagy in synovium and PBMCs of HLA-B27 positive patients with ankylosing spondylitis. Response to: 'Evidence that autophagy, but not the unfolded protein response, regulates the expression of IL-23 in the gut of patients with ankylosing spondylitis and subclinical gut inflammation' by Ciccia *et al.* *Annals of the rheumatic diseases* **73**, e68, doi:10.1136/annrheumdis-2014-206277 (2014).
- 14 Melis, L. *et al.* Systemic levels of IL-23 are strongly associated with disease activity in rheumatoid arthritis but not spondyloarthritis. *Annals of the rheumatic diseases* **69**, 618-623, doi:10.1136/ard.2009.107649 (2010).
- 15 Artis, D. & Spits, H. The biology of innate lymphoid cells. *Nature* **517**, 293-301, doi:10.1038/nature14189 (2015).
- 16 Glatigny, S. *et al.* Proinflammatory Th17 cells are expanded and induced by dendritic cells in spondylarthritis-prone HLA-B27-transgenic rats. *Arthritis and rheumatism* **64**, 110-120, doi:10.1002/art.33321 (2012).
- 17 Abe, Y. *et al.* Ankylosing enthesitis associated with up-regulated IFN-gamma and IL-17 production in (BXSB x NZB) F(1) male mice: a new mouse model. *Modern rheumatology / the Japan Rheumatism Association* **19**, 316-322, doi:10.1007/s10165-009-0166-0 (2009).
- 18 Jansen, D. T. *et al.* IL-17-producing CD4+ T cells are increased in early, active axial spondyloarthritis including patients without imaging abnormalities. *Rheumatology (Oxford, England)* **54**, 728-735, doi:10.1093/rheumatology/keu382 (2015).
- 19 Shen, H., Goodall, J. C. & Hill Gaston, J. S. Frequency and phenotype of peripheral blood Th17 cells in ankylosing spondylitis and rheumatoid arthritis. *Arthritis and rheumatism* **60**, 1647-1656, doi:10.1002/art.24568 (2009).
- 20 Jandus, C. *et al.* Increased numbers of circulating polyfunctional Th17 memory cells in patients with seronegative spondylarthritides. *Arthritis and rheumatism* **58**, 2307-2317, doi:10.1002/art.23655 (2008).
- 21 Xueyi, L. *et al.* Levels of circulating Th17 cells and regulatory T cells in ankylosing spondylitis patients with an inadequate response to anti-TNF-alpha therapy. *Journal of clinical immunology* **33**, 151-161, doi:10.1007/s10875-012-9774-0 (2013).
- 22 Kenna, T. J. *et al.* Enrichment of circulating interleukin-17-secreting interleukin-23 receptor-positive gamma/delta T cells in patients with active ankylosing spondylitis. *Arthritis and rheumatism* **64**, 1420-1429, doi:10.1002/art.33507 (2012).

- 23 Bautista-Caro, M. B. *et al.* Decreased Th17 and Th1 cells in the peripheral blood of patients with early non-radiographic axial spondyloarthritis: a marker of disease activity in HLA-B27(+) patients. *Rheumatology (Oxford, England)* **52**, 352-362, doi:10.1093/rheumatology/kes267 (2013).
- 24 Appel, H. *et al.* Analysis of IL-17(+) cells in facet joints of patients with spondyloarthritis suggests that the innate immune pathway might be of greater relevance than the Th17-mediated adaptive immune response. *Arthritis research & therapy* **13**, R95, doi:10.1186/ar3370 (2011).
- 25 Benham, H. *et al.* Interleukin-23 mediates the intestinal response to microbial beta-1,3-glucan and the development of spondyloarthritis pathology in SKG mice. *Arthritis & rheumatology (Hoboken, N.J.)* **66**, 1755-1767, doi:10.1002/art.38638 (2014).
- 26 Armaka, M. *et al.* Mesenchymal cell targeting by TNF as a common pathogenic principle in chronic inflammatory joint and intestinal diseases. *The Journal of experimental medicine* **205**, 331-337, doi:10.1084/jem.20070906 (2008).
- 27 Elewaut, D. *et al.* Enrichment of T cells carrying beta7 integrins in inflamed synovial tissue from patients with early spondyloarthropathy, compared to rheumatoid arthritis. *The Journal of rheumatology* **25**, 1932-1937 (1998).
- 28 Mackley, E. C. *et al.* CCR7-dependent trafficking of RORgamma(+) ILCs creates a unique microenvironment within mucosal draining lymph nodes. *Nature communications* **6**, 5862, doi:10.1038/ncomms6862 (2015).
- 29 Mielants, H., Veys, E. M., Cuvelier, C., De Vos, M. & Botelberghe, L. HLA-B27 related arthritis and bowel inflammation. Part 2. Ileocolonoscopy and bowel histology in patients with HLA-B27 related arthritis. *The Journal of rheumatology* **12**, 294-298 (1985).
- 30 Van Praet, L. *et al.* Microscopic gut inflammation in axial spondyloarthritis: a multiparametric predictive model. *Annals of the rheumatic diseases* **72**, 414-417, doi:10.1136/annrheumdis-2012-202135 (2013).
- 31 Van Praet, L. *et al.* Degree of bone marrow oedema in sacroiliac joints of patients with axial spondyloarthritis is linked to gut inflammation and male sex: results from the GIANT cohort. *Annals of the rheumatic diseases* **73**, 1186-1189, doi:10.1136/annrheumdis-2013-203854 (2014).